

REPORT DOCUMENTATION PAGE			Form Approved OMB No. 0704-0188	
Public reporting burden for this collection of information is estimated to average 1 hour per response, including the time for reviewing instructions, searching existing data sources, gathering and maintaining the data needed, and completing and reviewing the collection of information. Send comments regarding this burden estimate or any other aspect of this collection of information, including suggestions for reducing this burden, to Washington Headquarters Services, Directorate for Information Operations and Reports, 1215 Jefferson Davis Highway, Suite 1204, Arlington, VA 22202-4302, and to the Office of Management and Budget, Paperwork Reduction Project (0704-0188), Washington, DC 20503.				
1. AGENCY USE ONLY (Leave blank)		2. REPORT DATE December 2007		3. REPORT TYPE AND DATES COVERED Journal Article-Human Factors
4. TITLE AND SUBTITLE Thermal Comfort and Sensation in Men Wearing a Cooling System Controlled by Skin Temperature			5. FUNDING NUMBERS	
6. AUTHOR(S) C.R. Vernieuw, L.A. Stephenson, M.A. Kolka				
7. PERFORMING ORGANIZATION NAME(S) AND ADDRESS(ES) Thermal and Mountain Medicine Division U.S. Army Research Institute of Environmental Medicine Natick, MA 01760-5007			8. PERFORMING ORGANIZATION REPORT NUMBER M06-23	
9. SPONSORING / MONITORING AGENCY NAME(S) AND ADDRESS(ES) Same as #7 above			10. SPONSORING / MONITORING AGENCY REPORT NUMBER	
11. SUPPLEMENTARY NOTES				
12a. DISTRIBUTION / AVAILABILITY STATEMENT Approved for public release; distribution unlimited			12b. DISTRIBUTION CODE	
13. ABSTRACT (Maximum 200 words) The study was done to determine whether thermal comfort (TC), thermal sensation (TS), and subjective factors gauging environmental stress were negatively affected with different cooling methods in men exercising in chemical protective clothing. Previous studies have reported that intermittent regional cooling improved the efficacy of cooling as compared with constant cooling (CC), but no studies have addressed whether there is any improvement in thermal comfort. Eight male volunteers exercised at moderate work intensity (425 W) in three microclimate cooling tests. The circulating fluid in the cooling garment was provided during exercise to the head (6% body surface area [BSA]), torso (22% BSA), and thighs (44% BSA) and manipulated under three methods: (a) CC, (b) pulsed cooling (PC), and (c) PC activated by mean skin temperature (Tsk) control (PCskin). TC and TS ratings were recorded every 20 min during the 80-min test. TC and TS ratings were not different for PCskin and CC; thus the participants perceived PCskin and CC ( $p < .001$ ). In PCskin, Tsk was significantly higher than in PC and CC ( $p < .001$ ), and PCskin was rated as being not as warm as PC according to TS. This indicates that the PCskin method was perceived as being as cool as CC and cooler than PC. These findings indicate that the PCskin cooling method is an acceptable alternative to CC and PC based on human perceptions.				
14. SUBJECT TERMS skin temperature, thermal comfort, thermal sensation			15. NUMBER OF PAGES 12	
			16. PRICE CODE	
17. SECURITY CLASSIFICATION OF REPORT Unclassified	18. SECURITY CLASSIFICATION OF THIS PAGE Unclassified	19. SECURITY CLASSIFICATION OF ABSTRACT Unclassified	20. LIMITATION OF ABSTRACT Unclassified	

Working postures and physical  
Özok & G. Salvendy (Eds.), *Ad-*  
pp. 941-944). West Lafayette, IN:

*Evaluation of paramedic's tasks*  
of musculoskeletal injury (CUPE  
33). Richmond, British Columbia,  
of British Columbia.  
un, R. W. (1990). Estimating low  
endants using a hybrid anatomical  
Annual Conference of the Human  
p. 191-195). Mississauga, Ontario,  
tion of Canada.

an, R. W., Frank, J., Shannon, H.,  
ire and load sampling approach to  
in occupational settings. *Interna-*  
*nomics*, 27, 65-77.

*ropometry, ergonomics and design*  
or & Francis.

v. M., Herrin, G. D., & Chaffin, D.  
neutral trunk postures of autono-

*Journal of Work, Environment*  
human factors analysis of ambu-  
Eberts (Eds.), *Trends in ergonom-*  
*ings of the Second Mid-Central*  
sterdam: North-Holland.  
ring concerns in ambulance inter-  
In *Proceedings of the Human*  
ting (pp. 345-348). Santa Monica,  
mics Society.

of industrial engineering in  
ineering and Management  
tute of Technology. He re-  
ics and biomechanics from  
8.

nt in the Faculty of Archi-  
el Institute of Technology,  
n industrial design in 2004.

2006  
27

## Thermal Comfort and Sensation in Men Wearing a Cooling System Controlled by Skin Temperature

Carrie R. Vernieuw, Lou A. Stephenson, and Margaret A. Kolka, U.S. Army Research Institute of Environmental Medicine, Natick, Massachusetts

**Objective:** The study was done to determine whether thermal comfort (TC), thermal sensation (TS), and subjective factors gauging environmental stress were negatively affected with different cooling methods in men exercising in chemical protective clothing. **Background:** Previous studies have reported that intermittent regional cooling improved the efficacy of cooling as compared with constant cooling (CC), but no studies have addressed whether there is any improvement in thermal comfort. **Methods:** Eight male volunteers exercised at moderate work intensity (425 W) in three microclimate cooling tests. The circulating fluid in the cooling garment was provided during exercise to the head (6% body surface area [BSA]), torso (22% BSA), and thighs (44% BSA) and manipulated under three methods: (a) CC, (b) pulsed cooling (PC), and (c) PC activated by mean skin temperature ( $\bar{T}_{sk}$ ) control (PC<sub>skin</sub>). TC and TS ratings were recorded every 20 min during the 80-min test. **Results:** TC and TS ratings were not different for PC<sub>skin</sub> and CC; thus the participants perceived PC<sub>skin</sub> as being similar to CC. TS was significantly warmer with PC than with PC<sub>skin</sub> and CC ( $p < .001$ ). In PC<sub>skin</sub>,  $\bar{T}_{sk}$  was significantly higher than in PC and CC ( $p < .001$ ), and PC<sub>skin</sub> was rated as being not as warm as PC according to TS. **Conclusion:** This indicates that the PC<sub>skin</sub> method was perceived as being as cool as CC and cooler than PC. **Application:** These findings indicate that the PC<sub>skin</sub> cooling method is an acceptable alternative to CC and PC based on human perceptions.

### INTRODUCTION

Previously it was reported that intermittent regional cooling, as compared with a constant cooling paradigm, improved the efficacy of cooling exercising men wearing chemical protective clothing (Cheuvront, Kolka, Cadarette, Montain, & Sawka, 2003). Heat removal was significantly improved because mean skin temperature ( $\bar{T}_{sk}$ ) and, consequently, skin blood flow remained elevated for more time in the body regions supplied with regional cooling, having the effect of decreasing the insulation of the body surface and increasing radiative, convective, and conductive heat loss. Importantly, regional cooling was achieved with reduced circulation of the cooled fluid, presumably making dissipation of body heat more efficacious. The study did not address whether or not regional cooling resulted in a similar improvement in user comfort.

The rationale for the research was that any cooling paradigm that improved or maintained the user's cardiovascular and thermoregulatory integrity would improve the user's comfort when he or she exercised while wearing chemical protective clothing. Consequently, this study was initiated to further improve the cooling paradigm to optimize cooling and power conservation. The users' subjective ratings of how the new cooling paradigm made them feel relative to constantly supplied cooling were measured. In the current study, the participants' perception of thermal sensation and thermal comfort was determined to further support the effectiveness of pulsed cooling based on skin temperature feedback from subjective ratings by the user.

The physiological approach used in our laboratory to improve cooling efficacy focused on retaining a warm skin so that the cutaneous vasculature remains vasodilated at a level that supports rapid

convective and radiative heat loss once cool liquid circulates. This provided a microenvironment appropriate for heat transfer away from the body.

The research aim was to further improve efficiency in cooling by using pulsatile cooling controlled by temperature sensors placed on the skin ( $PC_{skin}$ ), so that real-time  $\bar{T}_{sk}$  could be used to activate the fluid-circulating pump when  $\bar{T}_{sk}$  increased  $1.5^{\circ}\text{C}$  above thermoneutral skin temperature. When the warm skin was cooled to a temperature of  $33.5^{\circ}\text{C}$  (slightly warmer than thermoneutral skin temperature), the pump was automatically deactivated. Kolka et al. (2004) documented the heat transfer properties of this most recent physiologic, integrative approach used in our laboratory to optimize microclimate cooling. In addition, Stephenson, Vernieuw, Leammukda, and Kolka (2007) described these findings in greater detail. In short, integrating the individual's  $\bar{T}_{sk}$  response to activate cooling was as effective as constant cooling and time-activated pulsed cooling to improve heat dissipation in men exercising in protective clothing. This improved efficacy occurred with a 46% reduction in electrical power used.

Thermal comfort (TC) was chosen as a measure because it was suspected that using the word "comfort" might help the participants integrate thermal sensation (TS) with the burden of exercise and wearing the cooling garment. Fanger (1970) defined TC as a "state in which he [the participant] expresses satisfaction with the thermal environment; i.e. he would prefer neither a warmer nor a colder environment" (p. 152). With this said, one would assume that a person's performance would be at his or her best when he or she is thermally comfortable (Fanger, 1970).

TS was used as another measurement for subjective perception to the cooling methods. TS provided specific descriptors for thermal perception from *very cold* to *very hot*, and this scale includes a reference to *neutral* TS (Berglund, 1998). It is important to point out that thermal balance, clothing, ambient temperature, ambient dew-point temperature, wind speed, radiant temperature, clothing, and metabolism all affect TS (Berglund, 1998). In the current study, the participants were not in thermal balance across the entire exercise bout, so one has to view TS and TC perception as a composite indicator for all the factors at the time of the measurement. The many factors that influence TC and TS were  $\bar{T}_{sk}$ , local skin temperature sites with and without cooling, core temperature, the liquid flow

rate in the cooling garment, the temperature of the liquid in the cooling garment at the skin, heart rate, skin wettedness, clothing, and metabolism.

The participants' perception of the entire experiment was measured using the Environmental Symptoms Questionnaire (ESQ), a measurement of subjective responses to extreme environments (Sampson, Kobrick, & Johnson, 1994). Embedded within the ESQ are five indices and nine factors. In the current study the most relevant indices were subjective heat illness, muscle discomfort, cardiopulmonary discomfort, tiredness, and well-being. The most relevant factors determined from the ESQ were distress, alertness, exertion, muscle discomfort, and fatigue.

The purpose of this research was to determine whether TC, TS, and subjective factors and indices from the ESQ were negatively affected in men exercising in chemical protective clothing when the cooling characteristics of fluid circulating through the cooling garment were changed. The research hypothesis was that the participants would perceive all cooling methods as equally cool.

## METHODOLOGY

The study was approved by institutional review boards based on scientific and human research review, and the investigators adhered to Army Regulation 70-25 and U.S. Army Research Institute of Environmental Medicine 70-25 on Use of Volunteers in Research. All participants were briefed regarding the risks and requirements to participate in the study. Participants who agreed to participate signed the consent form and then were medically cleared.

### Participants

Eight young, healthy, and fit men volunteered for this study: age  $20.6 \pm 1.7$  years, height  $175 \pm 0.05$  cm, weight  $73.14 \pm 7.45$  kg, body surface area (BSA)  $1.90 \pm 0.11$  m<sup>2</sup>, and body mass index (BMI)  $24.7 \pm 1.3$  (mean  $\pm$  SD). Women were not studied because none volunteered to participate.

### Experimental Procedures

Prior to testing, participants were thoroughly familiarized with all experimental techniques and fitted with the proper-sized liquid cooling garment, chemical protective clothing, and protective

mask. Participants were to testing. All testing Natick, Massachusetts explanation of TC and express their comfort provided during testir

Two participants were the morning (10:00) and (12:30), at the same time days to complete the different tests. Participants sensor (HQ, Inc., Palm instrument of core temperature to a data logger and ten Inc., Southborough, MA

After arriving at the drank 120 ml of water other fluid intake too seminude body mass electronic precision balance (Toledo ID1 Multirange Corp., New Berlin, WI) dressed in the inner clothing participants were instrumented with temperature/heat flow thermistors, electrodes, a heart rate monitor (CIC Inc., Port Washington, NY) and a temperature monitor. Participants wore the cooling garment, chemical protective clothing, and a biological field mask layer).

Body mass was recorded on, before the exercise ( $T_c$ ) and  $\bar{T}_{sk}$  were measured. Heart rate (HR) was monitored every 5 min using the recording system and checked the recording system (CIC, Switzerland) for safety. Measurements were recorded at 40, 60, and 80 min. Metabolism was measured by open-circuit spirometry (Vmax, Inc., Sandy, UT) during exercise for a 3- to 4-min period. The protective mask was complete, participants determined body mass seminude body mass.

**Clothing.** Participants wore the same clothing each trial. The person wore spandex shorts

t, the temperature of the  
nt at the skin, heart rate,  
, and metabolism.

tion of the entire exper-  
ng the Environmental  
(ESQ), a measurement  
extreme environments  
hinson, 1994). Embed-  
ve indices and nine fac-  
ie most relevant indices  
ss, muscle discomfort,  
ort, tiredness, and well-  
actors determined from  
rtness, exertion, muscle

earch was to determine  
ctive factors and indices  
vely affected in men ex-  
ctive clothing when the  
uid circulating through  
changed. The research  
ticipants would perceive  
ually cool.

## LOGY

l by institutional review  
and human research re-  
s adhered to Army Reg-  
rmy Research Institute  
ie 70-25 on Use of Vol-  
urtparticipants were briefed  
uirements to participate  
who agreed to partici-  
rm and then were med-

nd fit men volunteered  
1.7 years, height  $175 \pm$   
45 kg, body surface area  
body mass index (BMI)  
/omen were not studied  
l to participate.

## res

pants were thoroughly  
imental techniques and  
ed liquid cooling gar-  
clothing, and protective

mask. Participants were not heat acclimated prior to testing. All testing took place in October in Natick, Massachusetts. Participants were given an explanation of TC and TS and the proper way to express their comfort and sensation on the scale provided during testing.

Two participants were tested each week, one in the morning (10:00) and one in the early afternoon (12:30), at the same time on 3 consecutive testing days to complete the three methods for the different tests. Participants ingested a temperature pill sensor (HQ, Inc., Palmetto, FL) to allow measurement of core temperature, which was telemetered to a data logger and temperature monitor (FitSense, Inc., Southborough, MA).

After arriving at the laboratory, the participants drank 120 ml of water. During the experiment, no other fluid intake took place. Each participant's seminude body mass was determined using an electronic precision balance scale accurate to  $\pm 50$  g (Toledo ID1 Multirange, United Scale & Engineer Corp., New Berlin, WI). The participants then dressed in the inner clothing worn for the test. The participants were instrumented with skin temperature/heat flow thermistors, electrocardiogram electrodes, a heart rate monitor (Polar Watch, Polar CIC Inc., Port Washington, NY), and the core temperature monitor. Participants were then fitted with the cooling garment (middle layer of clothing), chemical protective clothing, and M-40 chemical-biological field mask with hood (outer clothing layer).

Body mass was recorded with all equipment on, before the exercise began. Core temperature ( $T_c$ ) and  $\bar{T}_{sk}$  were measured every minute. Heart rate (HR) was monitored continuously and recorded every 5 min using the Polar heart rate telemetry system and checked with an electrocardiography recording system (Cardiovit AT-6, Schiller AG, Switzerland) for safety purposes. TC and TS estimations were recorded every 20 min (Time 0, 20, 40, 60, and 80 min). Metabolic rate was measured by open-circuit spirometry (True Max 2400 Parvo-medics, Inc., Sandy, UT) during the last 5 min of exercise for a 3- to 4-min duration after removal of the protective mask and hood. Once exercise was complete, participants were weighed again to determine body mass with all equipment and the seminude body mass.

**Clothing.** Participants dressed the same for each trial. The personal layer of clothing consisted of spandex shorts, cotton socks, and athletic

shoes. The cooling layer was a three-piece liquid cooling garment that covered the head (hood), torso (vest), and legs (pants). The layer worn was composed of a chemical protective suit that included a charcoal-impregnated overgarment (top and bottom), cotton glove liners, butyl gloves, and a M-40 chemical-biological field mask with hood. This clothing configuration provided approximate insulative value and vapor permeability characteristics of 2.1 and 0.32, respectively, based on still air copper manikin studies (Kolka, Stephenson, & Gonzalez, 1994).

The liquid cooling garment design consisted of cotton or Nomex® aramid fabric, depending on skin region, woven or laminated around small-diameter Tygon® tubing (internal diameter 2.5 mm) divided into multiple parallel circuits. The tubing length for the suit was estimated to be ~108 m. Total BSA covered was estimated in a previous study to be 72% (head = 6%, torso = 22%, and legs = 44%) using a Cyberware® three-dimensional head and whole body scanner (Cyberware, Inc., Monterey, CA; Chevront et al., 2003).

The cooling garment was connected to a temperature-controlled recirculating water bath through inlet-outlet umbilical tubes exiting the garment at the waist (torso and legs) or collar (head). The total flow rate for constant cooling was 1.2 L/min, and inlet water temperature was 21.5°C. Pulsed cooling and  $PC_{skin}$  flow rates were also 1.2 L/min when the perfusate was circulating. The cooling system was run off an independent power supply, which allowed the measurement of voltage and current usage during each of the three cooling tests. Flow rates and inlet-outlet temperatures were also measured.

**Design.** All three tests were done in a warm, dry environment (dry bulb temperature = 30°C; dew-point temperature = 11°C, equivalent to 30% relative humidity). Participants walked on a treadmill for 80 min during each of the three experiments (1.36 m/s, 2% grade, ~225 W/m<sup>2</sup>; 425 W, wind speed 0.939 m/s). The circulating fluid for each region was controlled and manipulated under one of three methods for each test: (a) constant cooling (CC) provided continuous liquid cooling throughout exercise to all regions; (b) pulsed cooling was provided for 2 min, and then the circulating pump was deactivated for 2 min in alternating cycles (PC); (c) pulsed cooling was achieved by control of  $\bar{T}_{sk}$  in which cooling was activated at 34.5°C and deactivated when  $\bar{T}_{sk}$  decreased to 33.5°C ( $PC_{skin}$ ).



## Calculations

We calculated  $\bar{T}_{sk}$  from the formula (Stephenson et al., 2007)

$$\bar{T}_{sk} = 0.07T_{head} + 0.10T_{upper\ back} + 0.10T_{lower\ back} + 0.10T_{chest} + 0.10T_{abdomen} + 0.14T_{forearm} + 0.19T_{thigh} + 0.20T_{calf}$$

The subscripted type identifies the location where each temperature was measured.

TC and TS measurements were taken at 20-min intervals (Time 0, 20, 40, 60, and 80 min). The data sheet used for each was an open-ended magnitude estimation scale that was a 122-mm line with two labeled categories: *warm* and *cool* for TS and *discomfort* and *comfort* for TC on a scale adapted from Gagge, Stolwijk, and Saltin (1969; see Tables 1 and 2). The open-ended scale was devised to eliminate some of the constraints often imposed by a category scale (Marks, Borg, & Ljunggren, 1983). TS results were then identified on a 5-point scale of *cool* through *neutral* to *warm*. TC results were rated similarly, but on a 4-point scale of *comfortable* to *very uncomfortable* (Gagge et al., 1969).

Data were analyzed using commercial software (SigmaStat, SPSS Science, Chicago, IL). ANOVA (treatment by time) with repeated measures was performed on  $\bar{T}_{sk}$ , TC and TS, HR, power,  $T_{re}$ , cooling, and time of day. Tukey's HSD post hoc test was applied when significant or interaction events were found. When the test of normality failed with two-way repeated measures, which is often attributable to small sample size and the conservative nature of the test, the Holm-Sidak test was used. Statistical significance was set at  $p < .05$ .

## RESULTS

### Perception Effects for Cooling Method

In the current study, the participants did not perceive any difference in TC among CC, PC, and PC<sub>skin</sub> (Table 3). However, they discerned differences in TS among the cooling paradigms (Table 4,  $p \leq .001$ ). TS was rated significantly warmer in PC than in either CC or PC<sub>skin</sub> ( $p \leq .001$ ). There was no significant difference for TS between CC and PC<sub>skin</sub>. TS averaged *slightly warm* (approximately +5) for PC but was closer to *neutral* (+4) for CC and PC<sub>skin</sub>.

Figure 1 shows mean differences in TC and TS for the three cooling paradigms as a function of time of the experiment. The mean HR and  $\bar{T}_{sk}$  are also presented to show that there were no significant differences in HR, but the  $\bar{T}_{sk}$  averaged (means with SDs in parentheses) 32.33°C (0.89°C) during CC, 33.19°C (0.52°C) during PC, and 33.86°C (0.42°C) during PC<sub>skin</sub>, all significantly different from each other ( $p \leq .001$ ). This is particularly important because TS was rated higher in PC than in PC<sub>skin</sub>.

The ESQ responses were similar among the three cooling paradigms, further indicating that heat strain was perceived similarly among participants in all conditions (Table 5). This is verified by similar changes in core temperature and heart rate (Table 6). Table 5 shows the data for the indices and factors of interest for the ESQ in the current study. Note that the cooling methods did not discriminate among either negative or positive factors on the ESQ (Sampson et al., 1994).

### Perception Effects for Experimental Time

TC was significantly lower during the last

TABLE 1

Gagge

Warm

Slightly

Neutral

Slightly

Cool

minute of exercise (81 of the experiment ( $p =$  imately 2 (35.35%) or ble 3). At 75 min the h for measurement of v Although  $T_{re}$  was at it exercise, as compared 6),  $T_{head}$  decreased w ( $T_{head}$   $p \leq .001$  between 60 min vs. 80).

TS was greater at t ment, before walking ing was activated ( $T_{re}$  exercise time when cor 20, 40, 60, and 80 min of +5, or *slightly warm* lowest at 80 min and v or *slightly cool* (36.12 and mask were remov cantly different from t tal time ( $p < .05$ ). TS r were not significantly For these three exerci imated +4, or *neutral*

Figure 2 shows  $T_{re}$  through the pump (w ing pump activation ar ipants, P5 and P8. The anthropometric chara BSA = 1.79 m<sup>2</sup>, height ± 0.2 kg; P8: BMI = 24

TABLE 1: Thermal Comfort

Gagge et al., 1969	Numerical Code	Open-Ended Rating	Percentage
Very uncomfortable	4	Discomfort	100% 75%
Uncomfortable	3		60% 50%
Slightly uncomfortable	2		35% 25%
Comfortable	1	Comfort	10% 0%

## TS

## Cooling Method

e participants did not  
C among CC, PC, and  
they discerned differ-  
ling paradigms (Table  
significantly warmer in  
 $T_{skin}$  ( $p \leq .001$ ). There  
ce for TS between CC  
lightly warm (approx-  
closer to *neutral* (+4)

ferences in TC and TS  
ligms as a function of  
e mean HR and  $T_{sk}$  are  
at there were no sig-  
, but the  $T_{sk}$  averaged  
ases) 32.33°C (0.89°C)  
2°C) during PC, and  
C<sub>skin</sub>, all significantly  
≤ .001). This is partic-  
S was rated higher in

re similar among the  
further indicating that  
imilarly among partic-  
ble 5). This is verified  
temperature and heart  
ws the data for the in-  
for the ESQ in the cur-  
oling methods did not  
egative or positive fac-  
et al., 1994).

## Experimental Time

ower during the last

## Percentage

100%  
75%  
60%  
50%  
35%  
25%  
10%  
0%

TABLE 2: Thermal Sensation

Gagge et al., 1969	Numerical Code	Open-Ended Rating	Percentage
Warm	6	Warm	100%
			90%
			80%
Slightly warm	5		77.50%
			70%
			60%
Neutral	4		57.50%
			50%
			40%
Slightly cool	3		37.50%
			30%
			20%
Cool	2	Cool	17.50%
			10%
			0%

minute of exercise (80 min) than at any other time of the experiment ( $p = .007$ ) and averaged approximately 2 (35.35%) or *slightly uncomfortable* (Table 3). At 75 min the hood and mask were removed for measurement of volume of oxygen consumed. Although  $T_c$  was at its highest toward the end of exercise, as compared with all other times (Table 6),  $T_{head}$  decreased when the hood was removed ( $T_{head} p \leq .001$  between trials and Time 0, 20, 40, 60 min vs. 80).

TS was greater at the beginning of the experiment, before walking started and before the cooling was activated (Time 0), as compared with any exercise time when cooling was operational (Times 20, 40, 60, and 80 min), and approximated a rating of +5, or *slightly warm* ( $70.25\% \pm 1.07\%$ ). TS was lowest at 80 min and was rated approximately +3, or *slightly cool* ( $36.13\% \pm 4.92\%$ ), after the hood and mask were removed. This rating was significantly different from that at any other experimental time ( $p < .05$ ). TS ratings at 20, 40, and 60 min were not significantly different from one another. For these three exercise times the ratings approximated +4, or *neutral* (Table 4).

Figure 2 shows TS, TC,  $T_{sk}$ , and the flow through the pump (which demonstrates the cooling pump activation and inactivation) for 2 participants, P5 and P8. These participants had similar anthropometric characteristics (P5: BMI = 24.7, BSA = 1.79 m<sup>2</sup>, height = 1.68 m, weight = 69.8 ± 0.2 kg; P8: BMI = 24.9; BSA = 1.79 m<sup>2</sup>; height =

1.68 m, weight 70.3 ± 0.6 kg), yet their ratings of TS and TC differ considerably, as can be seen on inspection of Figure 2.

P5 perceived himself to be at a *neutral* TS while he was standing in the chamber waiting for the exercise to begin during CC and PC at Time 0. He was *slightly warm* prior to the start of exercise during PC<sub>skin</sub>. During exercise itself, he was *slightly cool* (Times 20 and 40) to *cool* (Times 60 and 80) during CC, *neutral* (Time 20) to *slightly cool* (Times 40, 60, and 80) during PC, and *slightly cool* (Times 20, 40, and 60) to *cool* (Time 80) during PC<sub>skin</sub>.

P8, however, perceived the conditions to be warmer. During CC he started off *slightly warm* (Time 0), and once cooling was activated he rated TS as *neutral* (Times 20, 40, 60, and 80). During PC he was *slightly warm* (Times 0, 40, and 60) until the mask came off at the end of exercise, when he felt *neutral*. During PC<sub>skin</sub> he was *slightly warm* (Time 0) to *neutral* (Time 20) and then *slightly cool* for the last 40 min of exercise. This is also evident when looking at PC<sub>skin</sub> cooling cycles. P8 cooled more quickly once cooling was activated. P8 thus had nine cooling cycles as compared with P5, who had five cooling cycles.

There was no significant difference in the TC, TS, HR, or  $T_{sk}$  results between the participants who were tested in the morning and those who were tested in the late morning. There was, however, a significant difference in  $T_c$  ( $p = .035$ ) between the

Continued on page 1040

TABLE 3: Mean (SD) Thermal Comfort


Time (min)	CC (%)	CC Numerical Code	PC (%)	PC Numerical Code	PC <sub>skin</sub> (%)	PC <sub>skin</sub> Numerical Code
0: Standing, no cooling	60.6 (19.7)	3 Uncomfortable	49.6 (16.7)	2 Slightly uncomfortable	50.9 (19.1)	3 Uncomfortable
20: Walking	45.7 (20.2)	2 Slightly uncomfortable	53.4 (15.1)	3 Uncomfortable	47.7 (15.0)	2 Slightly uncomfortable
40: Walking	47.6 (9.8)	2 Slightly uncomfortable	53.4 (13.6)	3 Uncomfortable	50.9 (16.5)	3 Uncomfortable
60: Walking	48.0 (13.0)	2 Slightly uncomfortable	55.3 (11.0)	3 Uncomfortable	48.6 (19.9)	2 Slightly uncomfortable
80: Mask off	35.2 (17.1)	2 Slightly uncomfortable	39.4 (19.0)	2 Slightly uncomfortable	31.4 (24.1)	2 Slightly uncomfortable

Note. Main effect by cooling method  $p = .265$ , ns; main effect by time 80 min  $< 0$ , 20, 40, 60 min,  $p < .050$ . CC = constant cooling, PC = pulsed cooling, PC<sub>skin</sub> = PC regulated by mean skin temperature.

TABLE 4: Mean (SD) Thermal Sensation

Time (min)	CC (%)	CC Numerical Code/TS	PC (%)	PC Numerical Code/TS	PC <sub>skin</sub> (%)	PC <sub>skin</sub> Numerical Code/TS
0: Standing, no cooling	71.1 (20.4)	5 Slightly warm	70.6 (20.2)	5 Slightly warm	69.1 (14.3)	5 Slightly warm
20: Walking	43.5 (14.9)	4 Neutral	60.9 (16.9)	5 Slightly warm	45.7 (22.8)	4 Neutral
40: Walking	43.0 (16.8)	4 Neutral	62.6 (16.0)	5 Slightly warm	51.6 (21.5)	4 Neutral
60: Walking	39.3 (18.8)	3 Slightly cool	64.9 (19.2)	5 Slightly warm	45.0 (21.1)	4 Neutral
80: Mask off	32.4 (18.3)	3 Slightly cool	41.7 (21.7)	4 Neutral	34.3 (30.6)	3 Slightly cool

Note. Main effect by cooling method, TS  $p < .001$ , PC  $> CC$ , PC<sub>skin</sub>: main effect for time,  $p < .001$  time 0  $> 40$ , 60, 80 min. CC = constant cooling, PC = pulsed cooling, PC<sub>skin</sub> = PC regulated by mean skin temperature, TS = thermal sensation.



60: Walking	39.3 (18.8)	3	Slightly cool	64.9 (19.2)	5	Slightly warm	45.0 (21.1)	4	Neutral
80: Mask off	32.4 (18.3)	3	Slightly cool	41.7 (21.7)	4	Neutral	34.3 (30.6)	3	Slightly cool

Note. Main effect by cooling method, TS  $p < .001$ , PC > CC, PC<sub>skin</sub>; main effect for time,  $p < .001$  time 0 > 40, 60, 80 min. CC = constant cooling, PC = pulsed cooling, PC<sub>skin</sub> = PC regulated by mean skin temperature, TS = thermal sensation.




Figure 1. Mean skin temperature ( $\bar{T}_{sk}$ ), core temperature ( $T_{co}$ ), heart rate (HR), thermal comfort, and thermal sensation for the three cooling methods (CC = constant cooling, PC = pulsed cooling, PC<sub>skin</sub> = PC regulated by mean skin temperature; see text for details). The top line for each panel shows  $\bar{T}_{sk}$ , the middle line shows HR, and the bottom line shows  $T_{co}$  ( $p < .05$  for TS: PC > CC > PC<sub>skin</sub>;  $p < .05$  for : CC < PC < PC<sub>skin</sub>).



The choice to measure  $\bar{T}_{sk}$  during the study could be a function of the nature of the experiment by design when  $T_c$  was walked. However,  $\bar{T}_{sk}$  is a more sensitive TC and TS ratings than  $T_c$  (Frank, Raja, Bulcaca, 1998). This study supports the focus of the

PC <sub>skin</sub>	P
7.0 (5.3)	.5
0	ns
0.8 (0.9)	.4
0.4 (0.5)	.3
1.5 (1.4)	.6
4.0 (3.7)	.8
0.2 (0.2)	.8
3.2 (0.6)	.4
0.1 (0.1)	.6
0.3 (0.2)	.6
0.4 (0.4)	.7

pulsed cooling, PC<sub>skin</sub> = PC

tors, which shows that all three to prevent the symptoms of heat stress in environment while wearing protective clothing. Observations of the indices ESQ, it was concluded based on skin temperature as an alternative method to be perceived by the participants.

As in a previous study, it was found that strain when no cooling was provided (Mont et al., 2003). In that study, heart rate increased by  $1.7 \pm 0.3^\circ\text{C}$ ,  $5^\circ\text{C}$ , and heart rate was higher than resting heart rate. These findings in men were similar to supplemental cooling during exercise in similar

temperature ( $\Delta T_c$ ).

PC <sub>skin</sub>
33.9 (0.2)*
0.5 (0.3)
31 (10)
46 (30)*
22 (18)*

cooling, PC =

conditions (Kolka et al., 1994). Clearly, the cooling paradigms used cooled effectively.

The novel paradigm used in this study, of PC based on skin temperature feedback, is essentially an individual prescription for cooling based on real-time  $\bar{T}_{sk}$  data. This concept is demonstrated in Figure 2, as the cooling cycles for PC<sub>skin</sub> were different between P5 and P8. P5 had only five cooling cycles and P8 had nine, in order to keep  $\bar{T}_{sk}$  below  $34.5^\circ\text{C}$  and above  $33.5^\circ\text{C}$ . Given that P5 felt slightly cool by 20 min of exercise, and remained slightly cool throughout exercise according to the TS data, the case can be made that P5 might have preferred to have the lower  $\bar{T}_{sk}$  limit of  $33.5^\circ\text{C}$  raised to a warmer temperature. He might have even preferred to have the upper limit raised to a temperature warmer than  $34.5^\circ\text{C}$ . Alternatively, he might have been more comfortable if the temperature of the liquid coolant had been raised to a higher temperature. However, P5's data in Figure 2 show that  $\bar{T}_{sk}$  was about  $1^\circ\text{C}$  warmer in PC<sub>skin</sub> than in PC.

Now, if one looks at the  $\bar{T}_{sk}$  in Figure 1 for the PC<sub>skin</sub> cooling paradigm, the TS data warrant exploring the possibility of increasing the liquid coolant temperature circulating in the cooling garment. It is possible that the pump could be deactivated at a temperature  $>33.5^\circ\text{C}$ . Although there was little difference in electrical power used (P5: 108 W; P8: 116 W), P8 had 23 W more heat extracted, or cooling, than did P5.

The effect of raising the liquid coolant temperature in the PC<sub>skin</sub> cooling paradigm is being explored by biophysical modeling based on the current study. If the modeling effort supports the idea that the liquid cooling temperature can be raised in the PC<sub>skin</sub> cooling paradigm, as deduced from the TS data in the current study, further power savings may be possible. CC and PC cooling paradigms also need to be further explored to determine the optimal coolant temperature and to apply our findings for individual considerations and preferences, as Nevins, Gonzalez, Nishi, and Gagge (1975) recommended earlier.

The choice to measure TS and TC perceptions during the study could be debated, because the nature of the experiment included changing  $\bar{T}_{sk}$  by design when  $T_c$  was also changed as the men walked. However,  $\bar{T}_{sk}$  has been shown in subjective TC and TS ratings to be weighted more heavily (Frank, Raja, Bulcao, & Goldstein, 1999), which supports the focus of this study and the important

findings from PC<sub>skin</sub>. There was a significant difference in  $T_c$  ( $p = .035$ ) between the morning and late morning trials, which was expected, given the circadian variation in  $T_c$ . The participants were controlled as whether they participated in the morning or the late morning session (Stephenson et al., 1984). The participants were kept fairly comfortable and, according to the ESQ, were not stressed.

In hindsight, to make the PC<sub>skin</sub> test more sensitive, more measurements of thermal perceptions could be taken during exercise. To avoid a potential limitation of the study, more participants might have been tested, although the power analysis done prior to the study showed we had an adequate number. TS and TC measurements in PC<sub>skin</sub> varied because the cooling was determined by skin temperature, so there were some instances when the cooling was on or off during data collection and thus was not consistent between the participants. Despite these drawbacks, the TC, TS, and ESQ measures provided the necessary information to answer the question of whether the men were as comfortable during exercise when cooling was activated less under the integrative approach of PC<sub>skin</sub>, as compared with CC and PC.

## CONCLUSIONS AND RECOMMENDATIONS

The current subjective findings on thermal perception, combined with those of Kolka et al. (2004) and Stephenson et al. (2007), provide strong evidence to support the development of a cooling garment that can be controlled by the individual's  $\bar{T}_{sk}$  to manage heat stress for longer periods of time for a given battery power. Commercial development of the prototype PC<sub>skin</sub> will provide better cooling and, thus, improved operational capability. Future analyses of the optimal liquid coolant temperature could lead to further improvement in power savings as compared with CC and PC and to improved individual perception of thermal comfort and sensation. Additional research should determine how a number of factors (e.g., exercise intensity, exercise duration, and hotter ambient temperature) might change subjective perception of the PC<sub>skin</sub> cooling paradigm.

## ACKNOWLEDGMENTS

We gratefully acknowledge the volunteers who  
Continued on page 1044






Figure 2. Mean skin temperature ( $\bar{T}_{sk}$ ), core temperature ( $T_c$ ), head temperature ( $T_{head}$ ), perfusate status, thermal comfort, and thermal sensation (TS) for the three cooling methods (CC = constant cooling, PC = pulsed cooling,  $PC_{skin}$  = PC regulated by mean skin temperature; see text for details) for 2 participants (P5 and P8). The top line for each




panel shows  $T_c$ , the middle panel shows  $T_{sk}$ , and the bottom panel shows  $T_{head}$  (in the TS ratings on the right axis).

## Ratings



## P 8

## TS Ratings



usate status, thermal com-  
= pulsed cooling, PC<sub>skin</sub> =  
P8). The top line for each

panel shows  $T_c$ , the middle line shows  $T_{head}$ , and the bottom line shows  $T_{sk}$ . The faint lines in the bottom 2 panels for both P5 and P8 (in the Slightly Warm to Warm sections) demonstrate the cooling pump activation and inactivation (right axis).



participated in this research; the medical support; the engineers, Walter Teal and Brad Laprise; and the technical expertise of Bruce Cadarette, Walida Leammukda, Sharon McBride, Donna Merullo, Shari Hallas, Leslie Levine, and Scott Robinson from the U.S. Army Research Institute of Environmental Medicine and Natick Soldier Systems Center.

The views, opinions and/or assertions contained in this publication are those of the authors and should not be construed as an official Department of the Army position, policy, or decision unless so designated by other documentation.

Human participants took part in these studies after giving their free and informed voluntary consent. Investigators adhered to AR 70-25 and USARMC Regulation 70-25 on the Use of Volunteers in Research. For protection of human participants, the investigators adhered to policies of applicable Federal Law CFR 46.

## REFERENCES

- Berglund, L. G. (1998). Comfort and humidity. *ASHRAE Journal*, 40(8), 35–41.
- Cheuvront, S. N., Kolka, M. A., Cadarette, B. S., Montain, S. J., & Sawka, M. N. (2003). Efficacy of intermittent, regional microclimate cooling. *Journal of Applied Physiology*, 94, 1841–1848.
- Fanger, P. O. (1970). Conditions for thermal comfort: Introduction of a general comfort equation. In J. D. Hardy, A. P. Gagge, & J. A. J. Stolwijk (Eds.), *Physiological and behavioral temperature regulation* (pp. 152–175). Springfield, IL: Charles C. Thomas.
- Frank, S. M., Raja, S. N., Bulcao, C. F., & Goldstein, D. S. (1999). Relative contribution of core and cutaneous temperatures to thermal comfort and autonomic responses in humans. *Journal of Applied Physiology*, 86, 1588–1593.
- Gagge, A. P., Stolwijk, J. A. J., & Saltin, B. (1969). Comfort and thermal sensations and associated physiological responses during exercise at various ambient temperatures. *Environmental Research*, 2, 209–229.
- Kolka, M. A., Cadarette, B. S., Cheuvront, S. N., Teal, W., Laprise, B., & Stephenson, L. A. (2004). Power savings during intermittent microclimate cooling (IRC) worn under personal protective equipment (PPE) [Abstract]. *Aviation, Space, and Environmental Medicine*, 75, B23.
- Kolka, M. A., Stephenson, L. A., & Gonzalez, R. R. (1994). Thermoregulation in women during uncompensable heat stress. *Journal of Thermal Biology*, 19, 315–321.
- Marks, L. E., Borg, G., & Ljunggren, G. (1983). Individual differences in perceived exertion assessed by two new methods. *Perception and Psychophysics*, 34, 280–288.
- Nevins, R. J., Gonzalez, R. R., Nishi, Y., N., & Gagge, A. P. (1975). Effect of changes in ambient temperature and level of humidity on comfort and thermal sensations. *ASHRAE Transactions*, 81(2), 169–182.
- Sampson, J. B., Kobrick, J. L., & Johnson, R. F. (1994). Measurement of subjective reactions to extreme environments: The environmental symptoms questionnaire. *Military Psychology*, 6, 215–233.
- Stephenson, L. A., Vernieuw, C. R., Leammukda, W., & Kolka, M. A. (2007). Skin temperature feedback optimizes microclimate cooling. *Aviation, Space, and Environmental Medicine*, 78, 377–382.
- Stephenson, L. A., Wenger, C. B., O'Donovan, B. H., & Nadel, E. R. (1984). Circadian rhythm in sweating and cutaneous blood flow. *American Journal of Physiology*, 246(3, Pt. 2), R321–R324.
- Carrie R. Vernieuw currently resides in Boise, Idaho. She received her M.S. in exercise and movement science from the University of Oregon in 2001.
- Lou A. Stephenson is a research physiologist who retired from the U.S. Army Research Institute of Environmental Medicine and now resides in Logansport, Indiana. She received her Ph.D. in human performance from Indiana University in 1981.
- Margaret A. Kolka is a research physiologist who retired from the U.S. Army Research Institute of Environmental Medicine and was chief of the Biophysics and Biomedical Modeling Division. She now resides in Kauai, Hawaii. She received her Ph.D. in human performance from Indiana University in 1980.

Date received: October 10, 2006

Date accepted: July 10, 2007

## Effects of Sp Natural Lang

Ludovic Le Bigot,  
National de la Re  
University, Rennes,

**Objective:** T  
and user recei  
dialogue. Ba  
based dialog  
ronments and  
and reception  
pants perfor  
tion system  
(typing and  
**Results:** Me  
more frequen  
duction and  
were longer  
longer, and th  
The producti  
load. They h  
the discourse  
language wo  
this combina

## INTRC

With the advent of information systems, jing themselves "talki text or speech. Rese: dialogue, or HCD, has of speech- versus tey on the user's perforr and dialogue manage

The goal of this stud and text-based mode action shape human-c a range of performar

## Issues for Modal C

The fields of appl and various (Wilkie,

Address correspondence to  
poitiers.fr. *HUMAN FACTS*  
© 2007, Human Factors a